

PRODUCT LIST

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
2,4-D DMA	480 g /L 2,4-D, selective broadleaf herbicide	SL	39.8	Rubber, plantation (coconut, <3-year; oil palm, < 2-year), paddy			●
	600 g /L 2,4-D, selective broadleaf herbicide	SL	49.8	Paddy, oil palm			●
	720 g /L 2,4-D DMA, selective broadleaf herbicide	SL	58.0%	Rubber, coconut, oil palm; paddy, cereals, sorghum			●
2,4-D DIMETHYLAMINE + DIETHANOLAMINE SALTS	625 g /L 2,4-D dual salt, selective broadleaf herbicide	SL	49.5%	Winter cereals, fallow, millet, pastures, sugar cane, lawn, right-of-way, industrial, wood plantations	●	●	
2,4-D ETHYL HEXYL ESTER	680 g /L 2,4-D ester, selective broadleaf herbicide	EC	59.2%	Cereals, pastures (non-legumes), right-of-way, industrial, sugar cane	●	●	
2,4-D IPA	450 g /L 2,4-D IPA, specifically for glyphosate IPA compatibility	SL	39.0%	Winter cereals, fallow, pastures, maize, sweet corn, sugar cane, sorghum, millet, pasture, right-of-way, industrial, lawn, wood plantations	●		
2,4-D + PICLORAM (both as TRIISOPROPANOLAMINE SALT)	300 g /L 2,4-D, 75 g/L PICLORAM, selective broadleaf herbicide	SL	25.3% + 6.3%	Winter cereals, sugar cane, sorghum, maize, pasture, right-of-way, industrial	●		
ACETOCHLOR	840 g /L Acetochlor, selective pre-emergence herbicide for the control of annual grasses and many broadleaf weeds	EC	75.0%	grain and silage maize.		●	
ACIFLUORFEN	224 g/L ACIFLUORFEN present as the sodium salt. Selective control of certain broadleaf weeds and grasses.	SL	19.1%	Soybeans, mung beans, green beans, peanuts	●		
ALACHLOR	480 g/L ALACHLOR selective pre-emergence herbicide for the control of certain grasses	EC	48.0%	Maize, sweetcorn, soy beans and various vegetable crops.		●	
AMETRYN	500 g/L AMETRYN systemic herbicide for pre- and post-emergence of grasses and broadleaf weeds control	SC	45.9%	Sugar cane, pineapples, bananas		●	
ATRAZINE	900 g/kg ATRAZINE for the control of annual weeds in various crops	WG	90.0%	Sorghum, maize, sweet corn, saccaline, broom millet, sugar cane, canola, lupins and other crops	●		
BROMOXYNIL	200 g/L BROMOXYNIL present as the N-OCTANOYL ESTER	EC	19.0%	Wheat, oats, barley, cereal rye and triticale (alone or undersown), linseed, clover and lucerne, pastures, turf, fallow, non-crop, roadsides and rights-of-way	●		
BROMOXYNIL + MCPA 2-ETHYL HEXYL ESTER	200 g/L BROMOXYNIL present as the OCTANOATE + 200 g/L MCPA present as the ETHYL HEXYL ESTER for broadleaf weeds control	EC	18.5% + 18.5%	Cereals, linseed and grass pastures	●		
BUTROXYDIM	250 g/kg BUTROXYDIM for the control and suppression of various broadleaf weeds and grasses	WG	25.0%	Chickpeas, lucerne, lentils, canola, peanuts, soybeans, cotton, sunflowers	●		
CARFENTRAZONE-ETHYL	400 g/L CARFENTRAZONE-ETHYL for the control of certain broadleaf weeds	EC	37.2%	Fallows, forest plantations, commercial industrial and public service areas, nuts, grapevines, pastures, turf	●		

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
CARFENTRAZONE-ETHYL	400 g/L CARFENTRAZONE-ETHYL for the control of certain broadleaf weeds	EC	37.2%	Fallows, forest plantations, commercial industrial and public service areas, nuts, grapevines, pastures, turf	●		
CHLORSULFURON	750 g/kg CHLORSULFURON selective herbicide for the control of annual ryegrass and certain broadleaf weeds	WG	75.0%	Wheat, barley, oats, cereal rye and triticale	●		
CLETHODIM	240 g/L CLETHODIM for the control of certain grass weeds in various crops	EC	25.0%	Beetroot, cabbage, canola, celery, chickpeas, cotton, faba beans, field peas, forestry, lentils, lettuce, lupins, mung beans, non-bearing fruit trees, onions, ornamentals, peanuts, potatoes and soybeans	●	●	●
CLODINAFOP-PROPARGYL + CLOQUINTOCET-MEXYL	240 g/L CLODINAFOP-PROPARGYL + 60g/L CLOQUINTOCET-MEXYL for the control of Wild Oats, Paradoxa Grass (Annual Phalaris), Canary grass and annual Ryegrass	EC	22.4% + 5.6%	Wheat	●		
CLOMAZONE	30 g/kg CLOMAZONE for the control of certain annual grasses	GR	3.0%	Rice	●		
CLOPYRALID	300 g/L CLOPYRALID present as the TRIISOPROPANOLAMINE SALT for the control of a wide range of broadleaf weeds	SL	25.8%	Wheat, barley, oats, triticale, canola, pastures, fallow land, forests and industrial situations	●		
CLOPYRALID (as potassium salt)	750 g/kg CLOPYRALID present as the POTASSIUM SALT for the control of various broadleaf weeds	SG	75.0%	Wheat, barley, oats, triticale, canola, pastures, forestry and fallow land	●		
CYHALOFOP-BUTYL	100 g/L CYHALOFOP-BUTYL for the control of perennial grasses	EC	10.1%	Paddy			●
DICAMBA	500 g/L DICAMBA as Dimethylamine Salt. For the control of certain broadleaf weeds	SL	42.0%	Winter cereals, pastures, conservation tillage, sugar cane, turf, rice and non-crop areas	●		
DIURON	500 g/L DIURON pre- and post-emergent selective herbicide	SC	42.0%	Rubber, oil palm, sugar cane			●
	900 g/L DIURON for selective weed control	WG	90.0%	Asparagus, bananas, cotton, sugar cane, lupins, summer, wheat, triticale, cereal, oats, non-crop areas	●		
DIQUAT	200 g/L DIQUAT present as DIQUAT DIBROMIDE MONOHYDRATE for the control of a wide range of broadleaf weeds	SL	16.7%	Soya beans, sugar cane, sunflower, mung bean, linseed, lupin, sorghum, rice, orchard, pasture, wheat, oats, lucerne, aquatic areas	●	●	
ETHOFUMESATE	500 g/L of ETHOFUMESATE selective herbicide for pre- and post-emergence control of grasses and broadleaf weed	SC	44.2%	Pastures, beets, turf		●	

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
FLUMETSULAM	800 g/kg FLUMETSULAM as pre- and post-emergence control of a range of broadleaf weeds	WG	70.0%	winter cereals, clover, lucerne, pastures, chickpeas, field peas, lentils, maize, peanuts, soy beans	●	●	
FLUMIOXAZIN	500 g/kg FLUMIOXAZIN for rapid knockdown and control of various grass and broadleaved weeds when mixed with certain glyphosate herbicides	WG	50.0%	(PRESOWING) Barley, chickpeas, cotton, lentils, maize, oats, sorghum, wheat. (RESIDUAL CONTROL OF WEEDS) Grapevines, pome fruit, stone fruit, citrus fruit, tree nuts, olives	●		
FLUROXYPYR MEPTYL ESTER	200 g/L FLUROXYPYR, post-emergent broadleaf herbicide	EC	20.5%	For the control of a wide range of broadleaf weeds in fallow, lucerne, maize, millets, pastures, poppies, sorghum, sugar cane, sweetcorn, winter cereals.	●		●
	333 g /L FLUROXYPYR, high loaded post emergent herbicide	EC	31.1%	Also, for the control of woody weeds in agricultural non-crop areas, commercial and industrial areas, forests, pastures and rights-of-way	●	●	
GLUFOSINATE AMMONIUM	60 g /L GLUFOSINATE AMMONIUM, non-selective knockdown herbicide	SL	5.66%	Rubber, cacao, coffee, citrus, paddy (pre-sowing), oil palm, raspberries, avocado, banana, guava, kiwifruits, mango, papaya, pineapple, olive, pome and stone fruits, vineyards, forest plantations, right-of-way, line-marking on sport grounds, and other non-agricultural areas.			●
	150 g /L GLUFOSINATE AMMONIUM, non-selective knockdown herbicide	SL	13.5%				●
	200 g /L GLUFOSINATE AMMONIUM, non-selective knockdown herbicide	SL	18.0%		●	●	IN PROG.
	280 g /L GLUFOSINATE AMMONIUM, non-selective knockdown herbicide	SL	24.5%				IN PROG.
GLUFOSINATE - P (NEW)	105 g /L GLUFOSINATE AMMONIUM-P, non-selective knockdown herbicide A better isomer, commonly known as L-ISOMER.	SL	10.0%		-	-	-
GLYPHOSATE IPA	110 g /L GLYPHOSATE IPA, non selective systemic herbicide	SL	10.2%	Rubber, coconut, coffee, oil palm, non-agricultural land			●
	220 g /L GLYPHOSATE IPA, non selective systemic herbicide	SL	20.0%				●
	345 g /L GLYPHOSATE IPA, non selective systemic herbicide	SL	29.8%	Rubber, coconut, coffee, oil palm, non-agricultural land			●
	355 g /L GLYPHOSATE IPA, non selective systemic herbicide	SL	30.3%	Pasture, fallow, cotton (with shielded sprayer, aquatic weed control, unwanted tree control	●	●	
	450 g /L GLYPHOSATE IPA, non selective systemic herbicide	SL	37.5%	Variable of fruits trees, vines, pasture, fallow, cotton (with shielded sprayer	●		
	500 g /L GLYPHOSATE IPA, non selective systemic herbicide	SL	40.7%	Pasture, fallow, cotton (with shielded sprayer, aquatic weed control	●	●	
GLYPHOSATE MONOAMMONIUM	680 g/L GLYPHOSATE present as the MONO-AMMONIUM SALT	SG	68.0%	Pasture, fallow, cotton (with shielded sprayer, pre- & post harvest management	●		

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
GLYPHOSATE POTASSIUM	540 g /L GLYPHOSATE present as potassium salt, non selective systemic herbicide	SL	39.7%	Pasture, fallow, cotton (with shielded sprayer, aquatic weed control	●	●	
GLYPHOSATE POTASSIUM + MONOETHANOLAMINE	600 g/L GLYPHOSATE present as the POTASSIUM and MONOETHANOLAMINE SALTS	SL	45.45%	Pasture, fallow, cotton (with shielded sprayer, pre- & post harvest management	●		
HALOSULFURON-METHYL	750 g/kg HALOSULFURON-METHYL for selective post-emergence control of nutgrass	WG	75.0%	Sugar cane, corn/maize, sorghum, turf, cotton	●		
HALOXYFOP	520 g/L HALOXYFOP present as the HALOXYFOP-R-METHYL ESTER	EC	44.0%	Canola, chickpeas, lentils, lupins, lucerne, clover pastures, cotton, soy beans, sunflowers	●	●	
HEXAZINONE	900 g/kg of HEXAZINONE use as selective herbicide for the control of weeds in various situations	WG	90.0%	Forestry, lucerne (established), non-crop areas		●	
IMAZAPIC	240 g/L IMAZAPIC for pre-emergence control of certain annual grass and broadleaf weeds	SL	21.8%	fallows, sugar cane	●		
	700 g/kg IMAZAPIC for pre-emergence control of certain annual grass and broadleaf weeds	WG	70.0%	fallows, sugar cane	●		
ISOXAFLUTOLE	750 g/kg ISOXAFLUTOLE for the control and suppression of various broadleaf weeds and grasses	WG	75.0%	Sugar cane, chickpeas	●		
MCPA 2ETHYL HEXYL ESTER	570 g/L MCPA 2EH broadacre cereal selective herbicide	EC	54.0%	Wheat, barley, oats, triticale, rye, grass pasture	●		
MCPA (2-ETHYL HEXYL ESTER) + DIFLUFENICAN	250 g/L MCPA present as the ETHYL HEXYL ESTER + 25 g/L DIFLUFENICAN for the control of broadleaf weeds	EC	24.0% + 2.48%	Winter cereals and clover	●		
MCPA + DICAMBA	340 g/L MCPA (present as the dimethylamine salt) + 80 g/L DICAMBA (present as the dimethylamine salt)	SL	28.85% + 6.78%	Winter cereals, pastures, turf and non-crop areas	●		
MCPA DMA	750 g /L MCPA DMA selective broadleaf herbicide	SL	63.8%	Wheat, barley, oats, triticale, rye, grass pasture, rice, sugar cane, turf	●	●	
MCPA POTASSIUM	400 g /L MCPA potassium salt for control sedges in rice, and various broadleaf weeds in cereals	SL	40.0%	Rice / paddy			●
MCPA + PICLORAM	420 g /L MCPA + 26 g /L PICLORAM broadacre herbicide	SL	34.6% + 2.2%	Wheat, barley, oat, triticale, linseed	●		
MCPB	385 g/L MCPB as the sodium salt For the control of broadleaf weeds	SL	32.5%	Pastures, linseed, clover		●	
MCPB + MCPA	375 g/L MCPB and 25 g/L MCPA, both as the sodium salts for the control broadleaf weeds	SC	31.9% + 2.13%	Pastures (grass/clover), cereals, peas and in white clover and grass seed crops		●	

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
METAMITRON	700 g/L METAMITRON selective weed control herbicide	SC	57.8%	Red beets, sugar beets, fodder beets, mangolds		●	
	700 g/kg METAMITRON selective weed control herbicide	WG	70.0%	Red beets, sugar beets, fodder beets, mangolds		●	
METRIBUZIN	750 g/kg METRIBUZIN selective herbicide	WG	75.0%	Cereals, pastures, vegetables	●	●	
METSULFURON METHYL	200 g/kg METSULFURON METHYL for the control of brush and broadleaf weeds	WG	20.0%	Rubber plantation; oil palm, paddy, non-crop land, right of way (railway, roadside, etc.), commercial & industrial areas.			●
	600 g/kg METSULFURON METHYL for the control of brush and broadleaf weeds	WG	60.0%	Native pastures, rights of way, commercial and industrial areas, Barley, cereal, rye, triticale, wheat	●	●	
OXYFLUORFEN	240 g/L OXYFLUORFEN for the selective control of certain broadleaf and grass weeds	EC	23.7%	Almonds, apples, apricots, grapevines, peaches, plums, pears, pecans, walnuts, pyrethrum, mango, papayas, broccoli, cabbages, cauliflower, forestry	●	●	
PRETILACHLOR	300 g/L PRETILACHLOR for the control of sedges and certain broadleaf weeds in paddy	EC	28.7%	Paddy			●
PROPANIL	364 g/L PROPANIL rice herbicide	EC	35.0%	Paddy			●
	364 g/L PROPANIL rice herbicide, non-flammable formulation	EC	33.55%		●		
	800 g/kg PROPANIL for the control of broadleaf weeds and sedges in paddy	WG	80.0%				●
PROPYZAMIDE	500 g/kg PROPYZAMIDE for selective control of certain grasses and broad leaf weeds	WG	50.0%	Lettuce, sports turf, home lawns, legume seed crops, pastures	●		
PROSULFOCARB	800 g/L PROSULFOCARB for the control of Annual Ryegrass (Lolium rigidum) and other grass and broadleaf weeds	EC	76.9%	Barley, wheat	●		
PROSULFOCARB + S-METOLACHLOR	800 g/L PROSULFOCARB + 120 g/L S-METOLACHLOR, for control of annual ryegrass (Lolium rigidum) and toad rush (Juncus bufonius) in barley and wheat.	EC	76.5% + 11.5%	Barley, wheat	●		
PYROXASULFONE	850 g/kg PYROXASULFONE for the control wide range of grasses	WG	85.0%	Wheat (except durum wheat), maize	●		
PYRAZOSULFURON-ETHYL	100 g/kg PYRAZOSULFURON-ETHYL for the control of broadleaf weeds and sedges	WP	10.0%	Paddy			●
PYRAZOSULFURON-ETHYL	100 g/kg PYRAZOSULFURON-ETHYL for the control of broadleaf weeds and sedges	WP	10.0%	Paddy			●
QUINCLORAC	220 g/kg QUINCLORAC for post-emergent control of broadleaf weeds and certain grasses	SC	21.9%	Paddy			●

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
QUIZALOFOP-P-ETHYL	100 g/L QUIZALOFOP-P-ETHYL for the control of certain grass weeds in broadleaf crops	EC	9.7%	Beetroot, cabbage, canola, carrots, cauliflower, chickpeas, clover, pastures, honey dew melons, lucerne, lupins, onions, potatoes, radish, tomatoes, sunflowers, soy beans	●		
S-METOLACHLOR	960 g/L S-METOLACHLOR broad-spectrum pre-emergent herbicide	EC	96.0%	Broccoli, brussels sprouts, cabbages, cauliflowers, canola, cotton, maize, sweet corn, soybeans, sunflower, spinach, sugar cane, sweet potatoes, wheat, barley, oats, triticale	●	●	
SIMAZINE	900 g/kg SIMAZINE for the control of weeds in various crops	WG	90.0%	Apples, pears, almonds, asparagus, berry fruit, canola, vineyards, roses, chickpeas, lupins, forestry	●		
SULFOSULFURON	750 g/kg SULFOSULFURON selective herbicide for the control of certain weeds in wheat and triticale	WG	75.0%	Wheat, triticale	●		
TERBUTHYLAZINE	500 g/L TERBUTHYLAZINE for the control of broadleaf and grass weeds in various crops	SC	50.0%	Forestry, lucerne, maize / sweetcorn, peas, industrial areas		●	
	875 g/kg TERBUTHYLAZINE for the control of weeds in various crops	WG	87.5%	Chickpeas, faba beans, field peas, lupins.	●		
	900 g/kg TERBUTHYLAZINE as a residual and knockdown herbicide for the control of grass and broadleaf weeds	WG	90.0%	Lucerne, orchards, maize/sweetcorn, peas, industrial areas		●	
TERBUTRYN + MCPA	275 g/L TERBUTRYN + 160 g/L MCPA present as the POTASSIUM SALT for the control seedling broadleaf weeds	SC	27.5% + 16.0%	Wheat, barley, some varieties of oats, pasture and sugar cane	●		
TERBUTHYLAZINE + HEXAZINONE	435 g/litre TERBUTHYLAZINE and 65 g/litre HEXAZINONE	WG	43.5% + 6.5%	Forestry, industrial / non-cropping areas		●	
TRALKOXYDIM	400 g/kg TRALKOXYDIM For the control of wild oats and annual ryegrass and suppression of annual phalaris	WG	40.0%	Wheat, barley, rye, triticale	●		
TRIASULFURON	750 g/kg TRIASULFURON or pre-plant control of certain grasses and post emergent control of Wild Radish	WG	75.0%	Wheat, oats, barleys, triticale	●		
TRIALATE	500 g/L TRIALLATE for the control of Wild Oats in various crops	EC	47.4%	Barley, wheat, triticale, chickpeas, faba beans, lupins, peas, linseed, canola, safflower	●		
TRIBENURON-METHYL	750 g/kg TRIBENURON-METHYL for the control of certain broadleaf weeds in fallows and pre-crop situations	WG	75.0%	Fallow, pre-crop conditions	●		

HERBICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
TRICLOPYR + PICLORAM + AMINOPYRALID	300 g/L TRICLOPYR as the butoxyethyl + 100 g/L PICLORAM and 8 g/L AMINOPYRALID as the amine salt for the control of brush weeds	EC	30.0% + 10.0% + 8.0%	Pastures		●	
TRICLOPYR BUTOXYETHYL ESTER	228 g/L TRICLOPYR broadleaf and brush control herbicide	EC	23.1%	For control of a range of environmental and noxious woody and herbaceous weeds for both agricultural and non-agricultural areas.			●
	600 g /L TRICLOPYR broadleaf and brush control herbicide	EC	50.0%		●	●	
	755 g /L TRICLOPYR broadleaf and brush control herbicide	EC	60.4%		●		
TRICLOPYR BUTOXYETHYL ESTER + PICLORAM	300 g/L TRICLOPYR + 100 g/L PICLORAM brush herbicide	EC	26.4% + 8.8%		●	●	
TRIFLURALIN	480 g/L TRIFLURALIN pre-emergence herbicide for the control of annual grasses and certain broad leaf weeds in certain horticultural and agricultural crops	EC	45.2%	Chickpeas, red kidney beans, faba beans, pigeon peas, lentil, navy beans, soybeans, cotton, lucerne, peanuts, canola, safflower, sugar cane, sunflower, wheat, barley, triticale, broccoli, cabbage, cauliflowers, tomatoes, brussels sprouts, carrots, orchard, vineyards	●	●	

FUNGICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
AZOXYSTROBIN	250 g/L AZOXYSTROBIN	SC	22.94%	For the control of various diseases of Grapes, Potatoes, Tomatoes, Cucurbits, Avocados, Mangoes, Passionfruit, Pyrethrum, Olives,, Carrot, Nursery Stock, Ornamentals, Cut Flower/Foliage and Rubus crop, etc.	●	●	●
	800 g/L AZOXYSTROBIN For the control of various diseases on wide range of crops	WG	80.0%	Avocados, grapes, mangoes, tomatoes, potatoes, cucurbits	●		
AZOXYSTROBIN + DIFENOCONAZOLE	200 g/L AZOXYSTROBIN + 125 g/L DIFENOCONAZOLE	SC	18.5% + 11.6%	Control various diseases of vegetable and field crops, ranging from okra, capsicums, cucurbits, leafy vegetables, paddy, oil palm, mangoes, black peppers			IN PROG.
AZOXYSTROBIN + EPOXINCONAZOLE	120 g/L AZOXYSTROBIN + 100 g/L EPOXINCONAZOLE	SC	11.2% + 9.5%	For the control of certain fungal disease of wheat and barley	●	●	
BENOMYL	500 g/L BENOMYL both curative and protective action fungicide	WP	50.0%	Papaya, rubber trees, French beans, peanuts, vegetables, pepper, ornamentals, mangoes, pineapples, bananas, melons, tobacco			●
BOSCALID + PYRACLOSTROBIN	252 g/kg BOSCALID + 128 g/kg PYRACLOSTROBIN fungicides for the control of various diseases	WG	25.2% + 12.8%	Apples, pears, citrus, grapes, onions, stone fruits, cherries		●	
CAPTAN	900 g/kg CAPTAN for the control of certain diseases on various crops	WG	90.0%	Apples, pears, grapes, stone fruits, strawberries, ornamentals, turf	●		
CHLOROTHALONIL	500 g/L CHLOROTHALONIL	SC	40.0%	For the control of certain fungal disease of bananas, tobacco, tomatoes			●
	900 g/kg CHLOROTHALONIL for the control of fungal diseases	WG	90.0%	Almonds, apricots, cherries, grapes, peaches, plums, bananas, broccolis, cabbages, carrots, capsicums, cucurbits, okras, onions, leeks, shallots, peanuts, tomatoes	●		
COPPER	190 g/L COPPER (Cu) present as TRIBASIC COPPER SULPHATE for the control of various disease on a range of crops	SC		Almonds, apples, peaches, apricots, avocados, citrus, cherries, plums, mangoes, vines, walnuts, bananas, beans, brassicas, capsicums, carrots, lettuce,, beets, onions, cucurbits	●		
CYPRODINIL + FLUDIOXONIL	375g/kg CYPRODINIL + 250g/kg FLUDIOXONIL for the control of Botrytis, Anthracnose and Rhizopus	WG	37.5% + 25.0%	Onions, grapes, strawberries		●	
DIFENOCONAZOLE	250 g/L DIFENOCONAZOLE	EC	24.8%	For the control of various diseases of potatoes, paddy, bananas, macadamias, vegetable crops includes carrots, tomatoes and cucurbits; fruits such as papaya, bananas, watermelon, mangoes.	●	●	●

FUNGICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
DIFENOCONAZOLE + PROPICONAZOLE	150 g/L DIFENOCONAZOLE + 150 g/L PROPICONAZOLE	EC	13.9% + 13.9%	For control disease of paddy			●
EPOXICONAZOLE	125 g/L EPOXICONAZOLE	SC	11.90%	For the control of certain fungal disease of wheat and barley	●	●	
	750 g/kg EPOXICONAZOLE For the control of certain fungal diseases	WG	75.0%	Wheat, barley	●		
FLUAZINAM	500 g/L FLUAZINAM in the form of a suspension concentrate	SC	39.7%	Grapes, tomatoes, potatoes, vegetables		●	
HEXACONAZOLE	50 g/L HEXACONAZOLE in the form of a suspension concentrate	SC	4.8%	Vegetables, fruits, ornamentals			●
IPRODIONE	500 g/L IPRODIONE fungicide for the control of disease in a range of crops	SC	42.7%	Asparagus, grapes, stone fruits, tomatoes, turf, ornamentals		●	
KRESOXIM-METHYL	500 g/kg KRESOXIM-METHYL fungicide	WG	50.0%	Grapes, ornamentals, cucumbers			●
MANCOZEB	750 g/kg MANCOZEB broad- spectrum fungicide for the control of certain fungus	WG	75.0%	Citrus, grapes, passion fruits, pome fruits, stone fruits, peas, beets, carrots, capsicum, cucurbits, lettuce, potatoes, onion, tomatoes, peanuts, chickpeas, beans	●	●	
	800 g/kg MANCOZEB broad- spectrum fungicide for the control of certain fungus	WG	80.0%				●
FLUTRIAFOL	250 g/L FLUTRIAFOL	SC	22.8%	For the control of certain fungal diseases on wheat, barley and canola.	●		
	500 g/L FLUTRIAFOL	SC	42.0%		●		
PENCYCURON	250 g/L PENCYCURON used to control diseases caused by Rhizoctonia solani	SC	23.2%	Paddy, cabbages, seedlings, turf			●
PHOSPHOROUS ACID	400 g/L PHOSPHOROUS ACID	SL		A systemic fungicide for the control of root rot in avocado, root and collar rot in citrus, root and crown rot in ornamentals & root and heart rot in pineapples caused by phytophthora fungi. KEN-FOS 600 also controls downy mildew in grapes, poppies and cucurbits			●
	600 g/L PHOSPHOROUS ACID	SL	40.0%		●		

FUNGICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
PROPICONAZOLE	250 g/L PROPICONAZOLE	EC	25.0%	For certain fungal diseases of bananas, peanuts, perennial ryegrass, pineapples, stone fruit, sugarcane, wheat and other crops	●		●
	500 g/L PROPICONAZOLE	EC	45.9%		●		
PROPINEB	400 g/kg PROPINEB offer protection to crops from fungal diseases	WP	40.0%	Tomatoes, sweet potatoes, watermelon, tobacco, oil palm, spinach, paddy, ornamentals, mangoes, chillies, papayas			●
	800 g/kg PROPINEB offer protection to crops from fungal diseases	WP	80.0%				●
PROTHIOCONAZOLE	250 g/L PROTHIOCONAZOLE broad-spectrum fungicide offer protection for a range of crops	EC		Wheat, barley, oats, corns, soyabeans, chickpeas, peanuts, cotton		●	
PROTHIOCONAZOLE + TEBUCONAZOLE	210g/L PROTHIOCONAZOLE + 210g/L TEBUCONAZOLE for the control of various diseases in crop	SC	18.75% + 18.75%	Barley, wheat, oats, canola	●		
SULPHUR	800 g/kg SULPHUR for the control of powdery mildew, rust and mites	WG	80.0%	Citrus, apples, grape, ornamentals, strawberries, tomatoes, pomefruits	●	●	
TEBUCONAZOLE	250 g/L TEBUCONAZOLE	SC	23.8%	For the control of various disease of bananas, peanuts, cereal crops, vegetables and other crops			●
	430 g/L TEBUCONAZOLE	SC	38.3%		●	●	
TRIFLOXYSTROBIN	500 g/kg TRIFLOXYSTROBIN for the control of black spot and powdery mildew, Sclerotinia	WG	50.0%	Squash, stone fruits, citrus, apples, pears		●	

INSECTICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
ABAMECTIN	18 g/L ABAMECTIN for the control of pest mites on various crops	EC	1.8%	Pears, apples, cotton, capsicums, citrus, tomatoes, strawberries, ornamental crops	●		●
ALPHA-CYPERMETHRIN	20 g/L ALPHA-CYPERMETHRIN synthetic pyrethroid	EC	2.27%	Control of insect pests including helopeltis on cacao; heliothis (Helicoverpa spp.) on various crops; red legged earth mite and blue oat mite on certain field crops and pastures and certain pests on fruit and vegetable crops			●
	100 g/L ALPHA-CYPERMETHRIN synthetic pyrethroid	EC	10.6%		●	●	
BIFENTHRIN	100 g/L BIFENTHRIN general purpose insecticide	EC	10.7%	Control various of pest in cotton, tomatoes, alfalfa, bananas, wheat, barley, field peas, sugarcane, grapes, citrus, oil palm, ornamentals, and various of vegetable crops, also use for control subterranean termites	●	●	
CARBOSULFAN	50 g/kg CARBOSULFAN for the control of hinoceros beetle	GR	5.0%	Oil palm			●
CHLORANTRANILIPROLE	200 g/L CHLORANTRANILIPROLE for the control of certain species of insect pests in a range of vegetable crops.	SC	20.0%	Control pests in wide range of vegetables such as cabbage, lettuce, broccoli spinach, sweetcorn, paddy, potato.	●	●	IN PROG.
	350 g/kg CHLORANTRANILIPROLE for the control of Lepidopteran species of insect pests on pulse crop and fruits	WG	35.0%	Chickpea, soybeans, mung bean, apple. Pears, avocados.	●	●	
CHLORFENAPYR + CHLORANTRANILIPROLE	220g/L CHLORFENAPYR + 110g/L CHLORANTRANILIPROLE with stomach and contact action	SC	20.0% + 10.0%	Cotton,, vegetables, fruit trees, ornamental	-	-	-
CYFLUTHRIN	50 g/L CYFLUTHRIN for the control of mango pulp weevil, leaf beetle, mango shoot borer and mosquito bugs	EC	5.0%	Cocoa, mangoes			●
CYPERMETHRIN	50 g/L CYPERMETHRIN synthetic pyrethroid	EC	5.5%	Commonly use in vegetable crops (includes cabbage, eggplants, okra, Chinese broccoli, chilli, etc.) oil palms, sweetcorns			●
	55 g/L CYPERMETHRIN synthetic pyrethroid in EW formulation	EW	5.5%				●
	150 g/L CYPERMETHRIN synthetic pyrethroid	EC	16.0%				●
	166 g/L CYPERMETHRIN synthetic pyrethroid in EW formulation	EW	16.0%				●
CYPERMETHRIN + DIMETHOATE	20 g/L CYPERMETHRIN + 200 g/L Dimethoate combination insecticide	EC	2.04% + 20.4%	Vegetable crops such as okra, cabbage, and cauliflower			●

INSECTICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMULATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
DIAFENTHIURON	25 g/L DIAFENTHIURON for the control of thrip, diamondback moth caterpillar, tobacco cutworm caterpillar	SC	25.0%	Chilies, cabbages			●
DIAZINON	600 g/L DIAZINON for the control of insect pests on a range of crops	EW	56.1%	Apples, pears, avocados, oranges, mandarins, strawberries, tomatoes, ornamentals, pastures, cereals		●	
DIMETHOATE	400 g/L DIMETHOATE for the control of various of pests on vegetables crops	EC	38.0%	Chilies, cucumbers, tomatoes, cabbages, French beans			●
DIMETHOATE + CYPERMETHRIN	200 g/L DIMETHOATE + 20 CYPERMETHRIN + for the control of insect pests on vegetable crops	EC	20.4% + 2.04%	Okras, cabbages, cauliflowers			●
DELTAMETHRIN	25 g/L DELTAMETHRIN for the control of pests on a wide range of crops	EC	2.8%	Grapes, papayas, jackfruits, cocoa, mangoes, strawberries, bananas, vegetable crops			●
EMAMECTIN BENZOATE	17 g/L EMAMECTIN BENZOATE insecticide	EC	2.2%	Cabbage, cauliflowers, sweetcorn, spinach			●
FIPRONIL	50 g/L FIPRONIL broad-spectrum insecticide	SC	4.95%	Control various of pests in cabbage, turf, lawn, paddy, oil palm and non-agricultural areas			●
	800 g/kg FIPRONIL for the control of insect pests on fruits, brassicas, and more	WG	80.0%	Bananas, brassicas, sugarcane	●		
FLONICAMID	500 g/kg FLONICAMID for the control of aphids & psyllids	WG	50.0%	Potatoes, wheat, cotton, apples, ornamentals		●	
IMIDACLOPRID	200 g/L IMIDACLOPRID sucking insect control insecticide	SC	18.3%	Control damaging pests in sugarcane, vegetable crops such as capsicum, cucurbits, eggplant, sweet potato, tomatoes, etc.	●	●	
	350 g/L IMIDACLOPRID sucking insect control insecticide	SC	30.2%	Also use for treating subterranean termites.	●	●	
	600g/L IMIDACLOPRID flowable seed dressing insecticide	SC	47.0	Cotton, maize, sweet corns, sunflower, sorghum, canola, pastures, cereals	●	●	
INDOXACARB	150 g/L INDOXACARB for the control of various insect pests on a wide range of crops	OD/SC	14.3%	Closed head cabbages, brussels sprouts, chickpeas, adzuki beans, soya beans, cotton.	●	●	●
LAMBDA-CYHALOTHRIN	25 g/L LAMBDA-CYHALOTHRIN synthetic pyrethroid	EC	2.8%	Control of certain insect pests in barley, wheat, various of vegetables, canola, citrus, alfalfa, onion bulbs, pasture, potatoes, sorghum.			●
	250 g/L LAMBDA-CYHALOTHRIN synthetic pyrethroid in CS formulation	CS	23.6%		●	●	

INSECTICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
METHOMYL	225 g/L METHOMYL for the control of insect pests on various crops	SL	22.5%	Apples, brassicas, legume seed crops, chickpeas, citrus, cotton, grapes, lentils, leafy vegetables, pastures, potatoes, pears, peaches, sorghum, soya beans, maize, sunflowers, tea tree, tobacco, barley, wheat, oat, tomatoes.	●		
PIRIMICARB	500 g/kg PIRIMICARB for controlling aphids	WG	50.0%	Vegetable crops, vegetables, ornamental, cotton	●		
PYMETROZINE	500 g/kg PYMETROZINE insecticides for the control of various insect pests	WG	50.0%	Vegetable crops, vegetables, ornamental, cotton			●
SPIROTETRAMAT	240 g/L SPIROTETRAMET systemic insecticide for the control of armoured scales, mealy bugs, thrips and others in a range of crops	SC	22.4%	Apples, blueberries, citrus, tomatoes, potatoes, grapes, pears	●	●	
THIAMETHOXAM	250g/kg THIAMETHOXAM systemic insecticide that effectively control sucking and chew insects	WP	25.0%	Rice, cotton, corn, soybeans, vegetables, fruit trees	-	-	-

FUMIGANTS

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
PHOSPHINE	330 g/L PHOSPHINE from ALUMINIUM PHOSPHIDE fumigation product for the control of insects of stored products	TAB / bag chain	33.0%	Storage of cereal grains, crop commodities, warehouse	●		
METHAM SODIUM	423 g/L 423 g/L METHAM present as the SODIUM SALT for pre-planting treatment for the control of soil-borne pests	SC	35.0%	Planting bed & rows, potting soil, lawn	●		

MOLLUSCICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
METALDEHYDE	1.5 g/kg METALDEHYDE for the control of snails & slugs	PELLET	0.15%	A wide range of vegetables, paddy, fruits, ornamental, greenhouses, nurseries,	●		
	50 g/kg METALDEHYDE for the control of snails & slugs	GR	5.0%	A wide range of vegetables, paddy, fruits, ornamental, greenhouses, nurseries,			●

TECHNICAL

ACTIVE INGREDIENT	ORIGIN
2,4-D 2-ETHYLHEXYL ESTER 96%	MALAYSIA
MCPA 2-ETHYLHEXYL ESTER 95%	MALAYSIA

PLANT GROWTH REGULATOR

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
ETHEPHON	480, 720 g/L & 900 g/L ETHEPHON for crop thinning, loosening or ripening in various crops and for accelerating boll opening, defoliation and pre- conditioning	SL	48.0%	Apples, grapes, cherries, mandarins, oranges, peaches, pineapples, sugar cane, tomatoes, cotton		•	
		EC	72.0%		•		
		SC	90.0%		•		
GIBBERELIC ACID	200 g/kg & 900 g/kg GIBBERELIC ACID for foliar spray application to certain varieties promote desirable harvest effects.	SG	20.0%	Citrus, grapes, prunes, pastures	•		
		SG	80.0%			•	
MEPIQUAT	38 g/L & 250 g/L MEPIQUAT present as the CHLORIDE for the management of fruiting and vegetative growth in cotton	AC	3.8%	Cotton	•		
		AC	25.0%		•		
THIDIAZURON	500 g/L THIDIAZURON A high quality flowable formulation for the defoliation of cotton prior to harvest	SC	50.0%	Cotton	•		

RODENTICIDES

ACTIVE INGREDIENT	DESCRIPTION	FORMU- LATION	W/W	TARGET CROPS / AREAS (NOT LIMITED TO):	REGISTERED IN		
					AU	NZ	MY
BRODIFACOUM	0.03 g/kg of BRODIFACOUM	BB	0.003%	Paddy, plantations			•
BROMADIOLONE	0.05 g/kg of BROMADIOLONE	BB	0.005%	Paddy, plantations, orchards			•

ADJUVANTS

PRODUCT	ACTIVE INGREDIENT	USES
ALCOHOL ALKOXYLATE	1000 g/L ALCOHOL ALKOXYLATE	Wetting agent reducing surface tension on plant waxy surface hence enabling solution to spread evenly on plant to maximise coverage.
ETHOXYLATED ALCOHOLS	1000 g/litre ETHOXYLATED ALCOHOLS	Spray mixture adjuvant for use with fungicides, insecticide, herbicides, plant growth regulants and foliar applied nutrients to improve spray deposition and adherence.
NONYL PHENOL ETHOXYLATED	50 g/L NONYL PHENOL ETHOXYLATE	Wetting agent reducing surface tension on plant waxy surface hence enabling solution to spread evenly on plant to maximise coverage.
PARAFFINIC OIL + NONYL PHENOL ETHOXYLATE, ALCOHOL ETHOXYLATE	582 g/L PARAFFINIC OIL + 240 g/L NONYL PHENOL ETHOXYLATE, ALCOHOL ETHOXYLATE	A specially formulated blend of paraffinic oil, penetrants and surfactants to enhance wetting, spreading and uptake of many agricultural chemicals.
SOYAL PHOSPHORUS + PROPIONIC ACID	350 g/L SOYAL PHOSPHORUS + 350 g/L PROPIONIC ACID	Surfactant, Penetrant, Acidifier, and Drift Retardant
POLYETHER MODIFIED POLYSILOXANE	1020 g/L POLYETHER MODIFIED POLYSILOXANE	Non-ionic surfactant for use with Glyphosate and certain brush weed herbicide to improve coverage, penetration and uptake. It is also can be used with insecticides, fungicides, and foliar nutrient

FOLIAR FERTILISERS

PRODUCT NAME	CONTENT (N-P-K)	USES
VERDE	11 - 8 - 6 + TE	Foliar fertilizer that can be used on wide range of crops. To Improves growth, flowering, yield, and quality, particularly under stress conditions.
VERDE BLOOM	5 - 10 - 12 + TE	Foliar fertilizer that can be used on wide range of crops to promote fruiting and flowering for the plants.

WE ALSO WELCOME ENQUIRIES ABOUT OEM FORMULATION TAILORED TO YOUR NEEDS

A large, light blue, stylized graphic resembling a leaf or a petal, composed of two overlapping shapes, serves as a background for the text.

KENSO

Together we grow

www.kensocorp.com
export@kenso.com.my